

# Recommendations for Pregnancies in Patients with Crigler-Najjar Syndrome

J. H. Paul Wilson · Maarten Sinaasappel ·  
Fred K. Lotgering · Janneke G. Langendonk

Received: 5 August 2011 / Revised: 24 February 2012 / Accepted: 12 March 2012 / Published online: 22 April 2012  
© SSIEM and Springer-Verlag Berlin Heidelberg 2012

**Abstract** During pregnancy, the developing foetus in mothers with Crigler-Najjar type 1 and 2 is exposed to raised levels of unconjugated bilirubin, with the risk of neurotoxicity. We describe two pregnancies in a patient with Crigler-Najjar type 2, who was carefully monitored prior to and during pregnancy and phototherapy adjusted to maintain serum bilirubin levels below 200  $\mu\text{mol/l}$  and the bilirubin/albumin molar ratio below 50%. Both pregnancies resulted in normal delivery of healthy infants who had normal neurological development. A review of all reported pregnancies in Crigler-Najjar patients and a set of recommendations are presented.

## Introduction

Crigler-Najjar syndromes type 1 (OMIM # 218800) and type 2 (OMIM # 606785) are rare disorders caused by homozygous or compound heterozygous mutations in the gene UGT1A1 and characterised by markedly raised levels of unconjugated bilirubin. Before the introduction of

phototherapy, many patients died of kernicterus in the neonatal period. Patients now can expect to survive into adulthood when treated with intensive daily phototherapy or, in the case of Crigler-Najjar type 2, with phenobarbitone with or without phototherapy. There have been very few reports of the effects of maternal unconjugated hyperbilirubinaemia on the foetus. The main concern is that the unconjugated bilirubin, which can cross the placental barrier, might cause foetal damage, especially foetal kernicterus.

In the Gunn rat, which is an animal model of the Crigler-Najjar disease, the jaundiced female rat has been reported to be less fertile than the non-jaundiced one (Davis and Yeary 1979). In addition, there seems to be an increased chance of foetal abnormalities in the jaundiced Gunn female rat (Davis et al. 1983). In humans, patients with acute and chronic liver disease also tend to be less fertile and to have an increased chance of abortion, or foetal abnormality and possibly kernicterus (Furhoff 1974; Roszkowski and Pizarek-Miedzinska 1968; Cotton et al. 1981). However, conclusions based on findings in acute and chronic hepatitis or cirrhosis cannot be extrapolated to unconjugated hyperbilirubinaemia without other features of liver disease.

At the time one of our patients expressed the desire to have children, four case reports had been published on pregnancies in patients with Crigler-Najjar disease, with five successful pregnancies (Cahill and McCarthy 1989; Taylor et al 1991; Smith and Baker 1994; Ito et al 2001). Three concerned patients with Crigler-Najjar type 2, one of whom continued to take phenobarbitone throughout the pregnancy. All were delivered of healthy infants at term. The fourth patient was diagnosed with Crigler-Najjar type 1, with total blood bilirubin levels between 357 and 458  $\mu\text{mol/l}$  throughout pregnancy (1 mg/dl is 17.1  $\mu\text{mol/l}$ ). She underwent caesarean section. The infant had a cord total

---

Communicated by: Robin Lachmann

---

Competing interests: None declared

---

J.H.P. Wilson · J.G. Langendonk (✉)  
Department of Internal Medicine, Internal medicine, metabolic  
diseases Office D416 Erasmus MC, PO Box 2040, 3000  
CA Rotterdam, The Netherlands  
e-mail: j.langendonk@erasmusmc.nl

M. Sinaasappel  
Department of Paediatrics, Erasmus MC, Rotterdam,  
The Netherlands

F.K. Lotgering  
Department of Obstetrics and Gynaecology, Radboud University  
Nijmegen Medical Centre, Nijmegen, The Netherlands

**Table 1** Recommendations for the management of pregnancy in patients with Crigler-Najjar disease types 1 and 2

1. Discussion of potential problems and genetic counselling before becoming pregnant
2. Administration of folic acid prior to and during pregnancy
3. Regular assay of serum bilirubin and albumin and adjustment of phototherapy to keep the serum bilirubin < 200  $\mu\text{mol/l}$  and the bilirubin/albumin molar ratio < 50%
4. If the patient is on Phenobarbital, reduce dosage to 50–60 mg/day
5. Prenatal care and delivery in a tertiary care centre, with facilities for rapid bilirubin and albumin assays and phototherapy for mother and infant
6. Avoidance of drugs that increase unbound, unconjugated bilirubin (Robertson et al. 1991; Strauss et al. 2006)
7. Neurological testing of the child after birth and during early childhood

bilirubin of 410  $\mu\text{mol/l}$ , unconjugated 217  $\mu\text{mol/l}$ . This high percentage of direct reacting bilirubin is unexpected, and as the total bilirubin level of the mother and the cord blood are so similar, it could represent an artefact. Initially there were no neurological abnormalities and the infant appeared to develop normally but he was found to have some spasticity at 9 months of age (Taylor et al. 1991). We drew up a set of recommendations (Table 1), which we presented to and discussed with the Crigler-Najjar patients and their families, emphasising the limited evidence on the effects of raised maternal bilirubin levels on the developing fetus.

We have managed two successful pregnancies in a Crigler-Najjar type 2 patient, according to our guidelines. Both children are developing normally, both physically and mentally. We reviewed all reported pregnancies in Crigler-Najjar patients discovered by searching PubMed and Embase using the terms Crigler-Najjar and pregnancy.

### Case Report

A woman with Crigler-Najjar disease type 2, who had been followed since childhood in our hospital, had maintained her bilirubin levels between 200 and 250  $\mu\text{mol/l}$  with 1–2 h of phototherapy (10 lamps of 100 W) per day. She did not use Phenobarbital because of the side effects. She was compound heterozygote for the mutations L15R and S191F in the UDT1A1 gene.


At the age of 27 years, she became pregnant. At 9 weeks gestation her serum bilirubin was 234  $\mu\text{mol/l}$  and albumin 43 g/l. Total bilirubin and direct bilirubin were determined using commercially available diazo-based colorimetric assays on a Roche Modular Analytics P 800-Module (Roche Diagnostics Nederland, Almere, The Netherlands). As during the course of her disease less than 5% of the total bilirubin was direct reacting, in the later phases of the pregnancy, only total bilirubin was measured. The phototherapy was increased to 3 h/day resulting in bilirubin levels between 165 and 205  $\mu\text{mol/l}$ . Her serum albumin

dropped to 34 g/l. Delivery was forceps-assisted after induction because of mild proteinuria at 39 weeks. At delivery, her bilirubin was 239  $\mu\text{mol/l}$ . The healthy boy weighed 3440 g and had an Apgar score of 9 at 5 min. Cord blood bilirubin levels were the same as the mother's. The mother's bilirubin rose after delivery to 278  $\mu\text{mol/l}$  and her albumin dropped to 30 g/l (Fig. 1). Directly after birth, her son's bilirubin was 177  $\mu\text{mol/l}$  with albumin 35 g/l; phototherapy was given to the child for 3 days, bilirubin dropped to 10  $\mu\text{mol/l}$  with unchanged albumin.

Two years later, she gave birth to a healthy girl at 39 weeks, weighing 3,150 g with an Apgar score of 10 at 5 min. Bilirubin and albumin levels were similar to the first pregnancy. Both children have developed normally without hyperbilirubinaemia or neurological abnormalities during a follow up of 11 and 9 years, respectively.

### Discussion

Since the second pregnancy of our patient, there have been five more case reports on the outcome of pregnancy in patients with Crigler-Najjar type 1 and 2, giving a total of four Crigler-Najjar type 1 patients and eight Crigler-Najjar type 2, and 17 deliveries. The reports are summarised in Table 2. As can be seen, no neurological damage has been detected in the offspring, with one exception. The one exception was a Crigler-Najjar type 1 patient reported by Taylor et al. (1991) who had previously had three first trimester therapeutic abortions performed for social reasons and a prior miscarriage at which time she had a bilirubin level of 460  $\mu\text{mol/l}$ . This patient is not reported to have been treated for the hyperbilirubinaemia. In all the other patients it has been possible to reduce maternal unconjugated bilirubin levels by increasing the intensity or duration of the phototherapy, in two instances in combination with fortnightly albumin infusions to increase bilirubin binding capacity or and in some Crigler-Najjar type 2 patients by continuing low dose phenobarbital treatment. As expected, maternal bilirubin levels were higher in Crigler-Najjar type 1, who also required much longer daily exposure to blue light – from 12 to 16 h.



**Fig. 1** Maternal serum bilirubin and albumin in µmol/l during pregnancy and after delivery. B = mean values in the 2-year period prior to pregnancy; numbers = duration of pregnancy in days; D = delivery; D + number = day after delivery

**Table 2** Reports of Crigler-Najjar and pregnancy in the literature

| Reference      | CN type | Maternal age (years) | Maternal bilirubin during pregnancy (µmol/l)/ | Maternal bilirubin at delivery (µmol/l)/ | Cord blood or 1st day bilirubin (µmol/l) | Maternal treatment | Mode of delivery | Newborn treatment           | Foetal outcome            |
|----------------|---------|----------------------|---|--|--|--------------------|------------------|-----------------------------|---------------------------|
| Taylor 1991    | NA/ 1   | 21                   | 363–458                                       | 420                                      | 410 CB                                   | None               | CS               | PT                          | Quadriplegic at 18 months |
| Gajdos 2006    | 1       | 28                   | 230–280 (400 before PT)                       | 242                                      | 222 CB                                   | PT albumin         | CS               | PT                          | Healthy                   |
| Hannam 2009    | 1       | 26                   | 200–380 (526 before PT)                       | 317                                      | 323 D1                                   | PT                 | CS               | PT + Phb                    | Healthy                   |
| Hannam 2009    | 1       | 22                   | 399–510                                       | 480                                      | 420 CB                                   | PT albumin         | Spontaneous      | PT + exchange transfusion   | Healthy                   |
| Cahill 1989    | 2       | 22                   | 178–183                                       | 104                                      | 121 D1                                   | Phb                | Induced          | PT                          | Healthy                   |
| Smith 1994     | 2       | 24                   | 91–164  | 150                                      | 130 CB                                   | None               | Spontaneous      | None                        | Healthy                   |
| Ito 2001       | 2       | < 34                 | 55–116  |  |  | PT + Phb           | CS               | None                        | Healthy                   |
|                |         |                      | 75–173  | 65–94                                    | 87 CB                                    | PT + Phb           | CS               | None                        | Healthy                   |
| Holstein 2005  | 2       | 36                   | 72–94   | NA                                       | NA                                       | PT + Phb           | CS               | None                        | Healthy                   |
| Holstein 2011  |         | 38                   | 72–94   | NA                                       | NA                                       | Phb                | CS               | None                        | Healthy                   |
|                |         | 41                   | 72–94   | NA                                       | NA                                       | Phb                | CS               | None                        | None                      |
| Saxena 2005    | 2       | 24                   | NA  | 139                                      | 174 D1                                   | None               | Spontaneous      | None                        | Healthy                   |
|                |         | 25                   | 130   |  |  | None               | Spontaneous      | Blood transfusion, PT + Phb | Healthy                   |
| Passuello 2009 | 2       | 37                   | 68–154  | 67                                       | 94 D1                                    | Phb                | Spontaneous      | None                        | Healthy                   |
| Arora 2009     | 2       | 26                   | 137–144                                       | 144                                      | 227 D1                                   | None               | Spontaneous      | PT                          | Healthy                   |
| Current case   | 2       | 27                   | 165–205                                       | 239                                      | 187 D1                                   | PT                 | Induced          | PT                          | Healthy                   |
|                |         | 29                   | 168–256                                       | 195                                      | 146 D1                                   | PT                 | Spontaneous      | None                        | Healthy                   |

Features of pregnancies in patients with Crigler-Najjar reported in the literature, *PT* phototherapy, *Phb* phenobarbital, *CS* caesarean sections, *NA* not available, *CB* cord blood, *D1* blood taken on first day after delivery. Bilirubin levels reported in mg/dl were converted to µmol/l by multiplying by 17.1. Note that in the report by Saxena (2005) cord blood levels showed a surprisingly high level of conjugated bilirubin, which might be an artefact

In general, low doses of phenobarbital have been given (25–100 mg/day).

Unconjugated bilirubin can cross the placenta, and this is indeed the normal route for excretion during pregnancy as the foetal liver is unable to conjugate or excrete bilirubin (Macias et al. 2009; Gajdos et al. 2006; McDonagh 2007; 2010). It is conceivable that phototherapy of the mother might reduce exposure of the foetal brain to a greater extent than reflected by serum bilirubin levels as the photoisomers of bilirubin are more water soluble and do not cross the blood–brain barrier or presumably the placenta to the same extent as unconjugated bilirubin (Mreihil 2010). At birth, foetal unconjugated bilirubin levels are in most reports roughly equivalent to those found in their mothers with Crigler-Najjar disease; however, phototherapy is usually stopped several hours before delivery, and circulating levels of photoisomers would be expected to be decreasing. However, serum levels were lower than the mother's in both infants reported here, which is compatible with a reduced clearance of photoisomers by the placenta.

At present it is impossible to state what concentration of unconjugated bilirubin is non-neurotoxic for the developing foetus, but it seems reasonable to try to keep the maternal (and therefore the foetal) unconjugated bilirubin concentrations below 200  $\mu\text{mol/l}$  and the molar ratio between unconjugated bilirubin and serum albumin below 50% (Gajdos et al. 2006; Strauss et al. 2006). Current national guidelines state that a bilirubin over 170  $\mu\text{mol/l}$  is an indication for phototherapy in term infants without risk factors for kernicterus and 70  $\mu\text{mol/l}$  in high-risk infants.

In most patients with Crigler-Najjar who attended antenatal clinics, it has been possible to achieve these levels by increasing phototherapy, and, if needed, by infusions of human albumin.

Admission to hospital to facilitate monitoring and treatment of both mother and infant is necessary – as can be seen by the frequent need for phototherapy of the neonate. Careful planning of the admission, including use of a written protocol and frequent visits by the metabolic diseases team to monitor and educate the medical and nursing staff of the obstetrics unit, is essential.

## Conclusions

Regular monitoring and adjustment of treatment to keep bilirubin levels below 200  $\mu\text{mol/l}$  and bilirubin/albumin molar ratios below 50% during pregnancy in patients with Crigler-Najjar syndrome can result in good foetal outcome.

## Conflict of Interest Statement

None.

## References

- Arora N, Choudhary S (2009) Pregnancy with Crigler-Najjar syndrome type II. *J Obstet Gynaecol* 29:242–244
- Cahill DJ, McCarthy CF (1989) Pregnancy and the Crigler-Najjar syndrome. *J Obstet Gynecol* 9:213
- Cotton BD, Brock BJ, Schiffrin BS (1981) Cirrhosis and fetal hyperbilirubinemia. *Obstet Gynecol* 57(6 suppl):25–27S
- Davis DR, Yeary RA (1979) Impaired fertility in the jaundiced female (Gunn) rat. *Lab Anim Sci* 29:739–743
- Davis DR, Yeary RA, Lee K (1983) Improved embryonic survival in the jaundiced female rat fed activated charcoal. *Pediatr Pharmacol* 3:79–85
- Furhoff AK (1974) Fate of children born to women with jaundice in pregnancy. *Arch Gynäk* 217:165–172
- Gajdos V, Petit F, Trioche P et al (2006) Successful pregnancy in a Crigler-Najjar type I patient treated by phototherapy and semimonthly albumin infusions. *Gastroenterology* 131:921–924
- Hannam S, Moriarty P, O'Reilly H et al (2009) Normal neurological outcome in two infants treated with exchange transfusions born to mothers with Crigler-Najjar Type 1 disorder. *Eur J Pediatr* 168:427–429
- Holstein A, Plaschke A, Lohse P, Egberts EH (2005) Successful photo- and phenobarbital therapy during pregnancy in a woman with Crigler-Najjar syndrome type II. *Scand J Gastroenterol* 40:1124–1126
- Holstein A, Bryan CS (2011) Three consecutive pregnancies in a woman with Crigler-Najjar syndrome type II with good maternal and neonatal outcomes. *Dig Liver Dis* 43:170
- Ito T, Katagiri C, Ikeno S, Takahashi H, Nagata N, Terakawa N (2001) Phenobarbital following phototherapy for Crigler-Najjar syndrome type II with good fetal outcome: a case report. *J Obstet Gynaecol Res* 27:33–35
- Macias RI, Marin JJ, Serrano MA (2009) Excretion of biliary compounds during intrauterine life. *World J Gastroenterol* 15:817–828
- McDonagh AF (2007) Movement of bilirubin and bilirubin conjugates across the placenta. *Pediatrics* 119:1032–1033
- McDonagh AF (2010) Controversies in bilirubin biochemistry and their clinical relevance. *Semin Fetal Neonatal Med* 15:141–147
- Mreihil K, McDonagh AF, Nakstad B, Hansen TW (2010) Early isomerization of bilirubin in phototherapy of neonatal jaundice. *Pediatr Res* 67:656–659
- Passuello V, Puhl AG, Wirth S et al (2009) Pregnancy outcome in maternal Crigler-Najjar syndrome type II: a case report and systematic review of the literature. *Fetal Diagn Ther* 26:121–126
- Robertson A, Karp W, Brodersen R (1991) Bilirubin displacing effect of drugs used in neonatology. *Acta Paediatr Scand* 80:1119–1127
- Roszkowski I, Pizarek-Miedzinska D (1968) Jaundice in pregnancy. II. Clinical course of pregnancy and delivery and condition of neonate. *Am J Obstet Gynecol* 101:500–503
- Saxena P, Arora R, Minocha B (2005) Crigler-Najjar syndrome with pregnancy. *J Obstet Gynecol India* 55:270–271
- Smith JF Jr, Baker JM (1994) Crigler-Najjar disease in pregnancy. *Obstet Gynecol* 84:670–672
- Strauss KA, Robinson DI, Vreman HK, Puffenberger EG, Hart G, Morton DH (2006) Management of hyperbilirubinemia and prevention of kernicterus in 20 patients with Crigler-Najjar disease. *Eur J Pediatr* 165(5):306–319
- Taylor WG, Walkinshaw SA, Farquharson RG, Fisk RA, Gilmore IT (1991) Pregnancy in Crigler-Najjar syndrome. *Br J Obstet Gynaecol* 98:1290–1291